

## Chapter 9

# Integration

### Antiderivative

A function  $F(x)$  is called an **antiderivative** (or *primitive*) of function  $f(x)$ , if

$$F'(x) = f(x)$$

**Computation:**

Guess and verify

**Example:** We want the antiderivative of  $f(x) = \ln(x)$ .

Guess:  $F(x) = x(\ln(x) - 1)$

Verify:  $F'(x) = (x(\ln(x) - 1))' =$   
 $= 1 \cdot (\ln(x) - 1) + x \cdot \frac{1}{x} = \ln(x)$

But also:  $F(x) = x(\ln(x) - 1) + 5$

### Antiderivative

The antiderivative is denoted by symbol

$$\int f(x) dx + c$$

and is also called the **indefinite integral** of function  $f$ . Number  $c$  is called **integration constant**.

Unfortunately, there are no “*recipes*” for computing antiderivatives (but tools one can try and which may help).

There are functions where antiderivatives cannot be expressed by means of elementary functions.

E.g., the antiderivative of  $\exp(-\frac{1}{2}x^2)$ .

## Basic Integrals

Integrals of some elementary functions:

$f(x)$	$\int f(x) dx$
0	$c$
$x^a$	$\frac{1}{a+1} \cdot x^{a+1} + c$
$e^x$	$e^x + c$
$\frac{1}{x}$	$\ln x  + c$
$\cos(x)$	$\sin(x) + c$
$\sin(x)$	$-\cos(x) + c$

(Table is created by exchanging the columns in our list of derivatives.)

## Integration Rules

### ► Summation rule

$$\int \alpha f(x) + \beta g(x) dx = \alpha \int f(x) dx + \beta \int g(x) dx$$

### ► Integration by parts

$$\int f \cdot g' dx = f \cdot g - \int f' \cdot g dx$$

### ► Integration by substitution

$$\int f(g(x)) \cdot g'(x) dx = \int f(z) dz$$

with  $z = g(x)$  and  $dz = g'(x) dx$

## Example – Summation Rule

Antiderivative of  $f(x) = 4x^3 - x^2 + 3x - 5$ .

$$\begin{aligned} \int f(x) dx &= \int 4x^3 - x^2 + 3x - 5 dx \\ &= 4 \int x^3 dx - \int x^2 dx + 3 \int x dx - 5 \int dx \\ &= 4 \frac{1}{4} x^4 - \frac{1}{3} x^3 + 3 \frac{1}{2} x^2 - 5x + c \\ &= x^4 - \frac{1}{3} x^3 + \frac{3}{2} x^2 - 5x + c \end{aligned}$$

## Example – Integration by Parts

Antiderivative of  $f(x) = x \cdot e^x$ .

$$\int \underbrace{x}_f \cdot \underbrace{e^x}_{g'} dx = \underbrace{x}_f \cdot \underbrace{e^x}_g - \int \underbrace{1}_{f'} \cdot \underbrace{e^x}_g dx = x \cdot e^x - e^x + c$$

$$\begin{aligned} f = x &\Rightarrow f' = 1 \\ g' = e^x &\Rightarrow g = e^x \end{aligned}$$

## Example – Integration by Parts

Antiderivative of  $f(x) = x^2 \cos(x)$ .

$$\int \underbrace{x^2}_f \cdot \underbrace{\cos(x)}_{g'} dx = \underbrace{x^2}_f \cdot \underbrace{\sin(x)}_g - \int \underbrace{2x}_{f'} \cdot \underbrace{\sin(x)}_g dx$$

Integration by parts of the second terms yields:

$$\begin{aligned} \int \underbrace{2x}_f \cdot \underbrace{\sin(x)}_{g'} dx &= \underbrace{2x}_f \cdot \underbrace{(-\cos(x))}_g - \int \underbrace{2}_{f'} \cdot \underbrace{(-\cos(x))}_g dx \\ &= -2x \cdot \cos(x) - 2 \cdot (-\sin(x)) + c \end{aligned}$$

Thus the antiderivative of  $f$  is given by

$$\int x^2 \cos(x) dx = x^2 \sin(x) + 2x \cos(x) - 2 \sin(x) + c$$

## Example – Integration by Substitution

Antiderivative of  $f(x) = 2x \cdot e^{x^2}$ .

$$\int \exp(\underbrace{x^2}_{g(x)}) \cdot \underbrace{2x}_{g'(x)} dx = \int \exp(z) dz = e^z + c = e^{x^2} + c$$

$$z = g(x) = x^2 \Rightarrow dz = g'(x) dx = 2x dx$$

## Integration Rules – Derivation

Integration by parts follows from the product rule for derivatives:

$$\begin{aligned} f(x) \cdot g(x) &= \int (f(x) \cdot g(x))' dx = \int (f'(x)g(x) + f(x)g'(x)) dx \\ &= \int f'(x)g(x) dx + \int f(x)g'(x) dx \end{aligned}$$

Integration by substitution follows from the chain rule:

Let  $F$  be an antiderivative of  $f$  and let  $z = g(x)$ . Then

$$\begin{aligned} \int f(z) dz &= F(z) = F(g(x)) = \int (F(g(x)))' dx \\ &= \int F'(g(x))g'(x) dx = \int f(g(x))g'(x) dx \end{aligned}$$

### Problem 9.1

Compute the antiderivatives of the following functions by means of integration by parts.

- (a)  $f(x) = 2x e^x$
- (b)  $f(x) = x^2 e^{-x}$
- (c)  $f(x) = x \ln(x)$
- (d)  $f(x) = x^3 \ln x$
- (e)  $f(x) = x (\ln(x))^2$
- (f)  $f(x) = x^2 \sin(x)$

### Problem 9.2

Compute the antiderivatives of the following functions by means of integration by substitution.

- (a)  $\int x e^{x^2} dx$
- (b)  $\int 2x \sqrt{x^2 + 6} dx$
- (c)  $\int \frac{x}{3x^2 + 4} dx$
- (d)  $\int x \sqrt{x+1} dx$
- (e)  $\int \frac{\ln(x)}{x} dx$

## Problem 9.3

Compute the antiderivatives of the following functions by means of integration by substitution.

(a)  $\int \frac{1}{x \ln x} dx$

(b)  $\int \sqrt{x^3 + 1} x^2 dx$


(c)  $\int \frac{x}{\sqrt{5 - x^2}} dx$

(d)  $\int \frac{x^2 - x + 1}{x - 3} dx$

(e)  $\int x(x - 8)^{\frac{1}{2}} dx$


## Area

Compute the areas of the given regions.



$$f(x) = 1$$


$$\text{Area: } A = 1$$



$$f(x) = \frac{1}{1+x^2}$$

Approximation  
by step function

## Riemann Sum



$$\xi_i = \frac{1}{2}(x_{i-1} + x_i)$$

in general:  $\xi_i \in (x_i, x_{i-1})$

$$A = \int_a^b f(x) dx \approx \sum_{i=1}^n f(\xi_i) \cdot (x_i - x_{i-1})$$

## Riemann Integral

$$I_n = \sum_{i=1}^n f(\xi_i) \cdot (x_i - x_{i-1})$$

is called a **Riemann sum** of  $f$ .

It can be shown that in many cases these Riemann sums converge when the length of the longest interval tends to 0.

This limit then is called the **Riemann integral** (or integral for short) of  $f$ .

## Riemann Integral – Properties

$$\int_a^b (\alpha f(x) + \beta g(x)) dx = \alpha \int_a^b f(x) dx + \beta \int_a^b g(x) dx$$

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\int_a^a f(x) dx = 0$$

$$\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$$

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx \quad \text{if } f(x) \leq g(x) \text{ for all } x \in [a, b]$$

## Fundamental Theorem of Calculus

Let  $F(x)$  be an antiderivative of a *continuous* function  $f(x)$ , then we find

$$\int_a^b f(x) dx = F(x) \Big|_a^b = F(b) - F(a)$$

By this theorem we can compute Riemann integrals by means of antiderivatives!

For that reason  $\int_a^b f(x) dx$  is called a **definite integral** of  $f$ .

### Example:

Compute the integral of  $f(x) = x^2$  over interval  $[0, 1]$ .

$$\int_0^1 x^2 dx = \frac{1}{3} x^3 \Big|_0^1 = \frac{1}{3} \cdot 1^3 - \frac{1}{3} \cdot 0^3 = \frac{1}{3}$$

## Integration Rules / (Definite Integrals)

### ► Summation rule

$$\int_a^b \alpha f(x) + \beta g(x) dx = \alpha \int_a^b f(x) dx + \beta \int_a^b g(x) dx$$

### ► Integration by parts

$$\int_a^b f \cdot g' dx = f \cdot g \Big|_a^b - \int_a^b f' \cdot g dx$$

### ► Integration by Substitution

$$\int_a^b f(g(x)) \cdot g'(x) dx = \int_{g(a)}^{g(b)} f(z) dz$$

with  $z = g(x)$  and  $dz = g'(x) dx$

## Example – Integration by Parts

Compute the definite integral  $\int_0^2 x \cdot e^x dx$ .

$$\begin{aligned} \int_0^2 \underbrace{x}_f \cdot \underbrace{e^x}_{g'} dx &= \underbrace{x}_f \cdot \underbrace{e^x}_g \Big|_0^2 - \int_0^2 \underbrace{1}_{f'} \cdot \underbrace{e^x}_g dx \\ &= x \cdot e^x \Big|_0^2 - e^x \Big|_0^2 = (2 \cdot e^2 - 0 \cdot e^0) - (e^2 - e^0) \\ &= e^2 + 1 \end{aligned}$$

Note: we also could use our indefinite integral from above,

$$\int_0^2 x \cdot e^x dx = (x \cdot e^x - e^x) \Big|_0^2 = (2 \cdot e^2 - e^2) - (0 \cdot e^0 - e^0) = e^2 + 1$$

## Example – Integration by Substitution

Compute the definite integral  $\int_e^{10} \frac{1}{\ln(x)} \cdot \frac{1}{x} dx$ .


$$\begin{aligned} \int_e^{10} \frac{1}{\ln(x)} \cdot \frac{1}{x} dx &= \int_1^{\ln(10)} \frac{1}{z} dz = \\ &= \ln(z) \Big|_1^{\ln(10)} = \\ &= \ln(\ln(10)) - \ln(1) \approx 0.834 \end{aligned}$$

$z = \ln(x) \Rightarrow dz = \frac{1}{x} dx$

## Example

Compute  $\int_{-2}^2 f(x) dx$  for function

$$f(x) = \begin{cases} 1 + x, & \text{for } -1 \leq x < 0, \\ 1 - x, & \text{for } 0 \leq x < 1, \\ 0, & \text{for } x < -1 \text{ and } x \geq 1. \end{cases}$$



We have

$$\begin{aligned} \int_{-2}^2 f(x) dx &= \int_{-2}^{-1} f(x) dx + \int_{-1}^0 f(x) dx + \int_0^1 f(x) dx + \int_1^2 f(x) dx \\ &= \int_{-2}^{-1} 0 dx + \int_{-1}^0 (1+x) dx + \int_0^1 (1-x) dx + \int_1^2 0 dx \\ &= \left(x + \frac{1}{2}x^2\right) \Big|_{-1}^0 + \left(x - \frac{1}{2}x^2\right) \Big|_0^1 \\ &= \frac{1}{2} + \frac{1}{2} = 1 \end{aligned}$$

## Problem 9.4

Compute the following definite integrals:

- (a)  $\int_1^4 2x^2 - 1 dx$
- (b)  $\int_0^2 3e^x dx$
- (c)  $\int_1^4 3x^2 + 4x dx$
- (d)  $\int_0^{\frac{\pi}{3}} \frac{-\sin(x)}{3} dx$
- (e)  $\int_0^1 \frac{3x+2}{3x^2+4x+1} dx$

## Problem 9.5

Compute the following definite integrals by means of antiderivatives:

- (a)  $\int_1^e \frac{\ln x}{x} dx$
- (b)  $\int_0^1 x(x^2+3)^4 dx$
- (c)  $\int_0^2 x\sqrt{4-x^2} dx$
- (d)  $\int_1^2 \frac{x}{x^2+1} dx$



### Problem 9.6

Compute the following definite integrals by means of antiderivatives:

(a)  $\int_0^2 x \exp\left(-\frac{x^2}{2}\right) dx$

(b)  $\int_0^3 (x-1)^2 x dx$

(c)  $\int_0^1 x \exp(x) dx$

(d)  $\int_0^2 x^2 \exp(x) dx$

(e)  $\int_1^2 x^2 \ln x dx$

### Problem 9.7

Compute  $\int_{-2}^2 x^2 f(x) dx$  for function

$$f(x) = \begin{cases} 1+x, & \text{for } -1 \leq x < 0, \\ 1-x, & \text{for } 0 \leq x < 1, \\ 0, & \text{for } x < -1 \text{ and } x \geq 1. \end{cases}$$

### Problem 9.8

Compute  $F(x) = \int_{-2}^x f(t) dt$  for function

$$f(x) = \begin{cases} 1+x, & \text{for } -1 \leq x < 0, \\ 1-x, & \text{for } 0 \leq x < 1, \\ 0, & \text{for } x < -1 \text{ and } x \geq 1. \end{cases}$$

## Summary

- ▶ antiderivate
- ▶ Riemann sum and Riemann integral
- ▶ indefinite and definite integral
- ▶ Fundamental Theorem of Calculus
- ▶ integration rules