


## Chapter 13

# Convex and Concave

# Monotone Functions\*

Function  $f$  is called **monotonically increasing**, if

$$x_1 \leq x_2 \Rightarrow f(x_1) \leq f(x_2)$$




It is called *strictly monotonically increasing*, if

$$x_1 < x_2 \Leftrightarrow f(x_1) < f(x_2)$$

Function  $f$  is called **monotonically decreasing**, if

$$x_1 \leq x_2 \Rightarrow f(x_1) \geq f(x_2)$$



It is called *strictly monotonically decreasing*, if

$$x_1 < x_2 \Leftrightarrow f(x_1) > f(x_2)$$

# Monotone Functions\*

For differentiable functions we have

$$\begin{aligned} f \text{ monotonically increasing} &\Leftrightarrow f'(x) \geq 0 \quad \text{for all } x \in D_f \\ f \text{ monotonically decreasing} &\Leftrightarrow f'(x) \leq 0 \quad \text{for all } x \in D_f \end{aligned}$$

$$\begin{aligned} f \text{ strictly monotonically increasing} &\Leftarrow f'(x) > 0 \quad \text{for all } x \in D_f \\ f \text{ strictly monotonically decreasing} &\Leftarrow f'(x) < 0 \quad \text{for all } x \in D_f \end{aligned}$$

Function  $f: (0, \infty), x \mapsto \ln(x)$  is strictly monotonically increasing, as

$$f'(x) = (\ln(x))' = \frac{1}{x} > 0 \quad \text{for all } x > 0$$

# Locally Monotone Functions\*

A function  $f$  can be monotonically increasing in some interval and decreasing in some other interval.

For *continuously* differentiable functions (i.e., when  $f'(x)$  is continuous) we can use the following procedure:

1. Compute first derivative  $f'(x)$ .
2. Determine all roots of  $f'(x)$ .
3. We thus obtain intervals where  $f'(x)$  does not change sign.
4. Select appropriate points  $x_i$  in each interval and determine the sign of  $f'(x_i)$ .

# Example – Locally Monotone Functions\*

In which region is function  $f(x) = 2x^3 - 12x^2 + 18x - 1$  monotonically increasing?

We have to solve inequality  $f'(x) \geq 0$ :

1.  $f'(x) = 6x^2 - 24x + 18$

2. Roots:  $x^2 - 4x + 3 = 0 \Rightarrow x_1 = 1, x_2 = 3$

3. Obtain 3 intervals:  $(-\infty, 1]$ ,  $[1, 3]$ , and  $[3, \infty)$

4. Sign of  $f'(x)$  at appropriate points in each interval:  
 $f'(0) = 3 > 0$ ,  $f'(2) = -1 < 0$ , and  $f'(4) = 3 > 0$ .

5.  $f'(x)$  cannot change sign in each interval:  
 $f'(x) \geq 0$  in  $(-\infty, 1]$  and  $[3, \infty)$ .

Function  $f(x)$  is monotonically increasing in  $(-\infty, 1]$  and in  $[3, \infty)$ .

# Monotone and Inverse Function

If  $f$  is *strictly monotonically increasing*, then

$$x_1 < x_2 \Leftrightarrow f(x_1) < f(x_2)$$

immediately implies

$$x_1 \neq x_2 \Leftrightarrow f(x_1) \neq f(x_2)$$

That is,  $f$  is *one-to-one*.




So if  $f$  is onto and strictly monotonically increasing (or decreasing), then  $f$  is **invertible**.

# Convex Set




A set  $D \subseteq \mathbb{R}^n$  is called **convex**, if for any two points  $\mathbf{x}, \mathbf{y} \in D$  the straight line segment between these points also belongs to  $D$ , i.e.,

$$(1 - h) \mathbf{x} + h \mathbf{y} \in D \quad \text{for all } h \in [0, 1], \text{ and } \mathbf{x}, \mathbf{y} \in D .$$

convex:




not convex:



# Intersection of Convex Sets

Let  $S_1, \dots, S_k$  be convex subsets of  $\mathbb{R}^n$ . Then their *intersection*  $S_1 \cap \dots \cap S_k$  is also convex.



The union of convex sets need not be convex.



# Example – Half-Space

Let  $\mathbf{p} \in \mathbb{R}^n$  and  $m \in \mathbb{R}$  be fixed,  $\mathbf{p} \neq 0$ . Then

$$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{p}^\top \cdot \mathbf{x} = m\}$$

is a so called **hyper-plane** which partitions the  $\mathbb{R}^n$  into two **half-spaces**

$$H_+ = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{p}^\top \cdot \mathbf{x} \geq m\} ,$$

$$H_- = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{p}^\top \cdot \mathbf{x} \leq m\} .$$

Sets  $H$ ,  $H_+$  and  $H_-$  are convex.

Let  $\mathbf{x}$  be a vector of goods,  $\mathbf{p}$  the vector of prices and  $m$  the budget.

Then the budget set is convex.

$$\{\mathbf{x} \in \mathbb{R}^n : \mathbf{p}^\top \cdot \mathbf{x} \leq m, \mathbf{x} \geq 0\}$$

$$= \{\mathbf{x} : \mathbf{p}^\top \cdot \mathbf{x} \leq m\} \cap \{\mathbf{x} : x_1 \geq 0\} \cap \dots \cap \{\mathbf{x} : x_n \geq 0\}$$


# Convex and Concave Functions

Function  $f$  is called **convex** in domain  $D \subseteq \mathbb{R}^n$ , if  $D$  is *convex* and


$$f((1 - h) \mathbf{x}_1 + h \mathbf{x}_2) \leq (1 - h) f(\mathbf{x}_1) + h f(\mathbf{x}_2)$$

for all  $\mathbf{x}_1, \mathbf{x}_2 \in D$  and all  $h \in [0, 1]$ . It is called **concave**, if

$$f((1 - h) \mathbf{x}_1 + h \mathbf{x}_2) \geq (1 - h) f(\mathbf{x}_1) + h f(\mathbf{x}_2)$$




convex



concave

# Concave Function\*

$$f((1-h)x_1 + hx_2) \geq (1-h)f(x_1) + hf(x_2)$$



Secant is below the graph of function  $f$ .

# Strictly Convex and Concave Functions

Function  $f$  is **strictly convex** in domain  $D \subseteq \mathbb{R}^n$ , if  $D$  is *convex* and

$$f((1-h)\mathbf{x}_1 + h\mathbf{x}_2) < (1-h)f(\mathbf{x}_1) + hf(\mathbf{x}_2)$$

for all  $\mathbf{x}_1, \mathbf{x}_2 \in D$  with  $\mathbf{x}_1 \neq \mathbf{x}_2$  and all  $h \in (0, 1)$ .

Function  $f$  is **strictly concave** in domain  $D \subseteq \mathbb{R}^n$ , if  $D$  is *convex* and

$$f((1-h)\mathbf{x}_1 + h\mathbf{x}_2) > (1-h)f(\mathbf{x}_1) + hf(\mathbf{x}_2)$$

for all  $\mathbf{x}_1, \mathbf{x}_2 \in D$  with  $\mathbf{x}_1 \neq \mathbf{x}_2$  and all  $h \in (0, 1)$ .

# Example – Linear Function

Let  $\mathbf{a} \in \mathbb{R}^n$  be fixed.

Then  $f(\mathbf{x}) = \mathbf{a}^\top \cdot \mathbf{x}$  is a linear map and we find:

$$\begin{aligned} f((1-h)\mathbf{x}_1 + h\mathbf{x}_2) &= \mathbf{a}^\top \cdot ((1-h)\mathbf{x}_1 + h\mathbf{x}_2) \\ &= (1-h)\mathbf{a}^\top \cdot \mathbf{x}_1 + h\mathbf{a}^\top \cdot \mathbf{x}_2 \\ &= (1-h)f(\mathbf{x}_1) + hf(\mathbf{x}_2) \end{aligned}$$

That is, every *linear function* is both *concave and convex*.

However, a linear function is neither strictly concave nor strictly convex, as the inequality is never strict.

# Example – Quadratic Univariate Function

Function  $f(x) = x^2$  is *strictly convex*:

$$\begin{aligned} & f((1-h)x + hy) - [(1-h)f(x) + hf(y)] \\ &= ((1-h)x + hy)^2 - [(1-h)x^2 + hy^2] \\ &= (1-h)^2 x^2 + 2(1-h)hxy + h^2 y^2 - (1-h)x^2 - hy^2 \\ &= -h(1-h)x^2 + 2(1-h)hxy - h(1-h)y^2 \\ &= -h(1-h)(x-y)^2 \\ &< 0 \quad \text{for } x \neq y \text{ and } 0 < h < 1. \end{aligned}$$

Thus

$$f((1-h)x + hy) < (1-h)f(x) + hf(y)$$

for all  $x \neq y$  and  $0 < h < 1$ ,

i.e.,  $f(x) = x^2$  is strictly convex, as claimed.

# Properties

- ▶ If  $f(\mathbf{x})$  is (strictly) *convex*, then  $-f(\mathbf{x})$  is (strictly) *concave* (and vice versa).
- ▶ If  $f_1(\mathbf{x}), \dots, f_k(\mathbf{x})$  are *convex* (concave) functions and  $\alpha_1, \dots, \alpha_k > 0$ , then

$$g(\mathbf{x}) = \alpha_1 f_1(\mathbf{x}) + \dots + \alpha_k f_k(\mathbf{x})$$

is also *convex* (concave).

- ▶ If (at least) one of the functions  $f_i(x)$  is *strictly convex* (strictly concave), then  $g(x)$  is strictly convex (strictly concave).


# Properties

For a differentiable functions the following holds:

- ▶ Function  $f$  is **concave** if and only if

$$f(\mathbf{x}) - f(\mathbf{x}_0) \leq \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)$$

i.e., the function graph is always below the tangent.




- ▶ Function  $f$  is **strictly concave** if and only if

$$f(\mathbf{x}) - f(\mathbf{x}_0) < \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) \quad \text{for all } \mathbf{x} \neq \mathbf{x}_0$$

- ▶ Function  $f$  is **convex** if and only if

$$f(\mathbf{x}) - f(\mathbf{x}_0) \geq \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)$$

(Analogous for strictly convex functions.)






# Univariate Functions\*

For two times differentiable functions we have

$$\begin{aligned} f \text{ convex} &\Leftrightarrow f''(x) \geq 0 && \text{for all } x \in D_f \\ f \text{ concave} &\Leftrightarrow f''(x) \leq 0 && \text{for all } x \in D_f \end{aligned}$$



Derivative  $f'(x)$  is  
monotonically decreasing,  
thus  $f''(x) \leq 0$ .

# Univariate Functions\*

For two times differentiable functions we have

$$f \text{ strictly convex} \iff f''(x) > 0 \quad \text{for all } x \in D_f$$

$$f \text{ strictly concave} \iff f''(x) < 0 \quad \text{for all } x \in D_f$$

# Example – Convex Function\*


Exponential function:

$$f(x) = e^x$$

$$f'(x) = e^x$$

$$f''(x) = e^x > 0 \quad \text{for all } x \in \mathbb{R}$$

$\exp(x)$  is (strictly) convex.



# Example – Concave Function\*


Logarithm function:  $(x > 0)$

$$f(x) = \ln(x)$$

$$f'(x) = \frac{1}{x}$$

$$f''(x) = -\frac{1}{x^2} < 0 \quad \text{for all } x > 0$$

$\ln(x)$  is (strictly) concave.



# Locally Convex Functions\*

A function  $f$  can be convex in some interval and concave in some other interval.

For two times *continuously* differentiable functions (i.e., when  $f''(x)$  is continuous) we can use the following procedure:

1. Compute second derivative  $f''(x)$ .
2. Determine all roots of  $f''(x)$ .
3. We thus obtain intervals where  $f''(x)$  does not change sign.
4. Select appropriate points  $x_i$  in each interval and determine the sign of  $f''(x_i)$ .

# Locally Concave Function\*

In which region is  $f(x) = 2x^3 - 12x^2 + 18x - 1$  concave?

We have to solve inequality  $f''(x) \leq 0$ .

1.  $f''(x) = 12x - 24$

2. Roots:  $12x - 24 = 0 \Rightarrow x = 2$

3. Obtain 2 intervals:  $(-\infty, 2]$  and  $[2, \infty)$

4. Sign of  $f''(x)$  at appropriate points in each interval:  
 $f''(0) = -24 < 0$  and  $f''(4) = 24 > 0$ .

5.  $f''(x)$  cannot change sign in each interval:  $f''(x) \leq 0$  in  $(-\infty, 2]$

Function  $f(x)$  is concave in  $(-\infty, 2]$ .

# Univariate Restrictions

Notice, that by definition a (multivariate) function is convex if and only if every restriction of its domain to a straight line results in a convex univariate function. That is:


Function  $f: D \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is convex

if and only if

$g(t) = f(\mathbf{x}_0 + t \cdot \mathbf{h})$  is convex

for all  $\mathbf{x}_0 \in D$  and

all non-zero  $\mathbf{h} \in \mathbb{R}^n$ .



# Quadratic Form

Let  $\mathbf{A}$  be a symmetric matrix  
and  $q_{\mathbf{A}}(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$  be the corresponding quadratic form.

Matrix  $\mathbf{A}$  can be diagonalized, i.e., if we use an orthonormal basis of its eigenvectors, then  $\mathbf{A}$  becomes a diagonal matrix with the eigenvalues of  $\mathbf{A}$  as its elements:

$$q_{\mathbf{A}}(\mathbf{x}) = \lambda_1 x_1^2 + \lambda_2 x_2^2 + \cdots + \lambda_n x_n^2 .$$

- ▶ It is convex if all eigenvalues  $\lambda_i \geq 0$   
as it is the sum of convex functions.
- ▶ It is concave if all  $\lambda_i \leq 0$   
as it is the negative of a convex function.
- ▶ It is neither convex nor concave if we have eigenvalues with  
 $\lambda_i > 0$  and  $\lambda_i < 0$ .



# Quadratic Form

We find for a quadratic form  $q_{\mathbf{A}}$ :

- ▶ *strictly convex*  $\Leftrightarrow$  *positive definite*
- ▶ *convex*  $\Leftrightarrow$  *positive semidefinite*
- ▶ *strictly concave*  $\Leftrightarrow$  *negative definite*
- ▶ *concave*  $\Leftrightarrow$  *negative semidefinite*
- ▶ *neither*  $\Leftrightarrow$  *indefinite*

We can determine the definiteness of  $\mathbf{A}$  by means of

- ▶ the eigenvalues of  $\mathbf{A}$ , or
- ▶ the (leading) principle minors of  $\mathbf{A}$ .

## Example – Quadratic Form

Let  $\mathbf{A} = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 3 & -1 \\ 0 & -1 & 2 \end{pmatrix}$ .      Leading principle minors:

$$A_1 = 2 > 0$$

$$A_2 = \begin{vmatrix} 2 & 1 \\ 1 & 3 \end{vmatrix} = 5 > 0$$

$$A_3 = |\mathbf{A}| = \begin{vmatrix} 2 & 1 & 0 \\ 1 & 3 & -1 \\ 0 & -1 & 2 \end{vmatrix} = 8 > 0$$

$\mathbf{A}$  is thus positive definite.

Quadratic form  $q_{\mathbf{A}}$  is *strictly convex*.

# Example – Quadratic Form

$$\text{Let } \mathbf{A} = \begin{pmatrix} -1 & 0 & 1 \\ 0 & -4 & 2 \\ 1 & 2 & -2 \end{pmatrix}. \quad \text{Principle Minors:}$$

$$A_1 = -1 \qquad A_2 = -4 \qquad A_3 = -2$$

$$A_{1,2} = \begin{vmatrix} -1 & 0 \\ 0 & -4 \end{vmatrix} = 4 \qquad A_{1,3} = \begin{vmatrix} -1 & 1 \\ 1 & -2 \end{vmatrix} = 1 \qquad A_{2,3} = \begin{vmatrix} -4 & 2 \\ 2 & -2 \end{vmatrix} = 4$$

$$A_{1,2,3} = \begin{vmatrix} -1 & 0 & 1 \\ 0 & -4 & 2 \\ 1 & 2 & -2 \end{vmatrix} = 0$$

$$\begin{array}{l} A_i \leq 0 \\ A_{i,j} \geq 0 \\ A_{1,2,3} \leq 0 \end{array}$$

$\mathbf{A}$  is thus negative semidefinite.

Quadratic form  $q_{\mathbf{A}}$  is *concave* (but not strictly concave).

# Concavity of Differentiable Functions

Let  $f: D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$  with Taylor series expansion

$$f(\mathbf{x}_0 + \mathbf{h}) = f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0) \cdot \mathbf{h} + \frac{1}{2} \mathbf{h}^\top \cdot \mathbf{H}_f(\mathbf{x}_0) \cdot \mathbf{h} + \mathcal{O}(\|\mathbf{h}\|^3)$$

*Hessian matrix*  $\mathbf{H}_f(\mathbf{x}_0)$  determines the concavity or convexity of  $f$  around expansion point  $\mathbf{x}_0$ .

- ▶  $\mathbf{H}_f(\mathbf{x}_0)$  *positive definite*  $\Rightarrow f$  *strictly convex* around  $\mathbf{x}_0$
- ▶  $\mathbf{H}_f(\mathbf{x}_0)$  *negative definite*  $\Rightarrow f$  *strictly concave* around  $\mathbf{x}_0$

- ▶  $\mathbf{H}_f(\mathbf{x})$  *positive semidefinite* for all  $\mathbf{x} \in D$   $\Leftrightarrow f$  *convex* in  $D$
- ▶  $\mathbf{H}_f(\mathbf{x})$  *negative semidefinite* for all  $\mathbf{x} \in D$   $\Leftrightarrow f$  *concave* in  $D$

# Recipe – Strictly Convex

1. Compute Hessian matrix

$$\mathbf{H}_f(\mathbf{x}) = \begin{pmatrix} f_{x_1x_1}(\mathbf{x}) & f_{x_1x_2}(\mathbf{x}) & \cdots & f_{x_1x_n}(\mathbf{x}) \\ f_{x_2x_1}(\mathbf{x}) & f_{x_2x_2}(\mathbf{x}) & \cdots & f_{x_2x_n}(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ f_{x_nx_1}(\mathbf{x}) & f_{x_nx_2}(\mathbf{x}) & \cdots & f_{x_nx_n}(\mathbf{x}) \end{pmatrix}$$

2. Compute all *leading principle minors*  $H_i$ .

3.

▶  $f$  strictly convex  $\Leftrightarrow$  all  $H_k > 0$  for (almost) **all**  $\mathbf{x} \in D$

▶  $f$  strictly concave  $\Leftrightarrow$  all  $(-1)^k H_k > 0$  for (almost) **all**  $\mathbf{x} \in D$

[  $(-1)^k H_k > 0$  implies:  $H_1, H_3, \dots < 0$  and  $H_2, H_4, \dots > 0$  ]

4. Otherwise  $f$  is *neither* **strictly** convex *nor* strictly concave.

# Recipe – Convex

1. Compute Hessian matrix

$$\mathbf{H}_f(\mathbf{x}) = \begin{pmatrix} f_{x_1x_1}(\mathbf{x}) & f_{x_1x_2}(\mathbf{x}) & \cdots & f_{x_1x_n}(\mathbf{x}) \\ f_{x_2x_1}(\mathbf{x}) & f_{x_2x_2}(\mathbf{x}) & \cdots & f_{x_2x_n}(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ f_{x_nx_1}(\mathbf{x}) & f_{x_nx_2}(\mathbf{x}) & \cdots & f_{x_nx_n}(\mathbf{x}) \end{pmatrix}$$

2. Compute all *principle minors*  $H_{i_1, \dots, i_k}$ .  
(Only required if  $\det(\mathbf{H}_f) = 0$ , see below)

3. ▶  $f$  convex  $\Leftrightarrow$  all  $H_{i_1, \dots, i_k} \geq 0$  for **all**  $\mathbf{x} \in D$ .

▶  $f$  concave  $\Leftrightarrow$  all  $(-1)^k H_{i_1, \dots, i_k} \geq 0$  for **all**  $\mathbf{x} \in D$ .

4. Otherwise  $f$  is *neither convex nor concave*.

# Recipe – Convex II

Computation of *all* principle minors can be avoided if  $\det(\mathbf{H}_f) \neq 0$ . Then a function is either strictly convex/concave (and thus convex/concave) or neither convex nor concave.

In particular we have the following recipe:

1. Compute Hessian matrix  $\mathbf{H}_f(\mathbf{x})$ .
2. Compute all *leading principle minors*  $H_i$ .
3. Check if  $\det(\mathbf{H}_f) \neq 0$ .
4. Check for strict convexity or concavity.
5. If  $\det(\mathbf{H}_f) \neq 0$  and  $f$  is neither strictly convex nor concave, then  $f$  is neither convex nor concave, either.

# Example – Strict Convexity

Is function  $f$  (strictly) concave or convex?

$$f(x, y) = x^4 + x^2 - 2xy + y^2$$

1. Hessian matrix:  $\mathbf{H}_f(\mathbf{x}) = \begin{pmatrix} 12x^2 + 2 & -2 \\ -2 & 2 \end{pmatrix}$

2. Leading principle minors:

$$H_1 = 12x^2 + 2 > 0$$

$$H_2 = |\mathbf{H}_f(\mathbf{x})| = 24x^2 > 0 \quad \text{for all } x \neq 0.$$

3. All leading principle minors  $> 0$  for almost all  $\mathbf{x}$   
 $\Rightarrow f$  is *strictly convex*. (and thus convex, too)



# Example – Cobb-Douglas Function

Let  $f(x, y) = x^\alpha y^\beta$  with  $\alpha, \beta \geq 0$  and  $\alpha + \beta \leq 1$ ,  
and  $D = \{(x, y) : x, y \geq 0\}$ .

Hessian matrix at  $\mathbf{x}$ :

$$\mathbf{H}_f(\mathbf{x}) = \begin{pmatrix} \alpha(\alpha - 1) x^{\alpha-2} y^\beta & \alpha\beta x^{\alpha-1} y^{\beta-1} \\ \alpha\beta x^{\alpha-1} y^{\beta-1} & \beta(\beta - 1) x^\alpha y^{\beta-2} \end{pmatrix}$$

Principle Minors:

$$H_1 = \underbrace{\alpha}_{\geq 0} \underbrace{(\alpha - 1)}_{\leq 0} \underbrace{x^{\alpha-2} y^\beta}_{\geq 0} \leq 0$$

$$H_2 = \underbrace{\beta}_{\geq 0} \underbrace{(\beta - 1)}_{\leq 0} \underbrace{x^\alpha y^{\beta-2}}_{\geq 0} \leq 0$$

# Example – Cobb-Douglas Function

$$\begin{aligned} H_{1,2} &= |\mathbf{H}_f(\mathbf{x})| \\ &= \alpha(\alpha - 1) x^{\alpha-2} y^\beta \cdot \beta(\beta - 1) x^\alpha y^{\beta-2} - (\alpha\beta x^{\alpha-1} y^{\beta-1})^2 \\ &= \alpha(\alpha - 1) \beta(\beta - 1) x^{2\alpha-2} y^{2\beta-2} - \alpha^2 \beta^2 x^{2\alpha-2} y^{2\beta-2} \\ &= \alpha\beta [(\alpha - 1)(\beta - 1) - \alpha\beta] x^{2\alpha-2} y^{2\beta-2} \\ &= \underbrace{\alpha\beta}_{\geq 0} \underbrace{(1 - \alpha - \beta)}_{\geq 0} \underbrace{x^{2\alpha-2} y^{2\beta-2}}_{\geq 0} \geq 0 \end{aligned}$$

$H_1 \leq 0$  and  $H_2 \leq 0$ , and  $H_{1,2} \geq 0$  for all  $(x, y) \in D$ .

$f(x, y)$  thus is *concave* in  $D$ .

For  $0 < \alpha, \beta < 1$  and  $\alpha + \beta < 1$  we even find:

$H_1 = H_1 < 0$  and  $H_2 = |\mathbf{H}_f(\mathbf{x})| > 0$  for almost all  $(x, y) \in D$ .

$f(x, y)$  is then *strictly concave*.

# Lower Level Sets of Convex Functions

Assume that  $f$  is *convex*.

Then the **lower level sets** of  $f$

$$\{\mathbf{x} \in D_f : f(\mathbf{x}) \leq c\}$$


are *convex*.

Let  $\mathbf{x}_1, \mathbf{x}_2 \in \{\mathbf{x} \in D_f : f(\mathbf{x}) \leq c\}$ ,  
i.e.,  $f(\mathbf{x}_1), f(\mathbf{x}_2) \leq c$ .

Then for  $\mathbf{y} = (1 - h)\mathbf{x}_1 + h\mathbf{x}_2$   
where  $h \in [0, 1]$  we find

$$\begin{aligned} f(\mathbf{y}) &= f((1 - h)\mathbf{x}_1 + h\mathbf{x}_2) \\ &\leq (1 - h)f(\mathbf{x}_1) + hf(\mathbf{x}_2) \\ &\leq (1 - h)c + hc = c \end{aligned}$$

That is,  $\mathbf{y} \in \{\mathbf{x} \in D_f : f(\mathbf{x}) \leq c\}$ , too.




# Upper Level Sets of Concave Functions

Assume that  $f$  is *concave*.


Then the **upper level sets** of  $f$

$$\{\mathbf{x} \in D_f : f(\mathbf{x}) \geq c\}$$

are *convex*.



upper level set



lower level set

# Extremum and Monotone Transformation

Let  $T: \mathbb{R} \rightarrow \mathbb{R}$  be a *strictly monotonically increasing* function.

If  $\mathbf{x}^*$  is a *maximum* of  $f$ , then  $\mathbf{x}^*$  is also a maximum of  $T \circ f$ .

As  $\mathbf{x}^*$  is a *maximum* of  $f$ , we have

$$f(\mathbf{x}^*) \geq f(\mathbf{x}) \text{ for all } \mathbf{x}.$$

As  $T$  is strictly monotonically increasing, we have

$$T(x_1) > T(x_2) \text{ falls } x_1 > x_2.$$

Thus we find

$$(T \circ f)(\mathbf{x}^*) = T(f(\mathbf{x}^*)) > T(f(\mathbf{x})) = (T \circ f)(\mathbf{x}) \text{ for all } \mathbf{x},$$

i.e.,  $\mathbf{x}^*$  is a maximum of  $T \circ f$ .


As  $T$  is one-to-one we also get the converse statement:

If  $\mathbf{x}^*$  is a *maximum* of  $T \circ f$ , then it also is a maximum of  $f$ .


# Extremum and Monotone Transformation

A strictly monotonically increasing Transformation  $T$  preserves the extrema of  $f$ .

Transformation  $T$  also preserves the level sets of  $f$ :



$$f(x, y) = -x^2 - y^2$$



$$T(f(x, y)) = \exp(-x^2 - y^2)$$

# Quasi-Convex and Quasi-Concave

Function  $f$  is called **quasi-convex** in  $D \subseteq \mathbb{R}^n$ , if  $D$  is *convex* and every *lower level set*  $\{\mathbf{x} \in D_f : f(\mathbf{x}) \leq c\}$  is *convex*.

Function  $f$  is called **quasi-concave** in  $D \subseteq \mathbb{R}^n$ , if  $D$  is *convex* and every *upper level set*  $\{\mathbf{x} \in D_f : f(\mathbf{x}) \geq c\}$  is *convex*.

# Convex and Quasi-Convex

Every *concave* (convex) function also is *quasi-concave* (and quasi-convex, resp.).

However, a quasi-concave function need not be concave.

Let  $T$  be a strictly monotonically increasing function.

If function  $f(\mathbf{x})$  is *concave* (convex), then  $T \circ f$  is *quasi-concave* (and quasi-convex, resp.).

Function  $g(x, y) = e^{-x^2 - y^2}$  is quasi-concave, as  $f(x, y) = -x^2 - y^2$  is concave and  $T(x) = e^x$  is strictly monotonically increasing.

However,  $g = T \circ f$  is not concave.



# A Weaker Condition

The notion of *quasi-convex* is **weaker** than that of *convex* in the sense that every convex function also is quasi-convex but not vice versa. There are much more quasi-convex functions than convex ones.

The importance of such a weaker notions is based on the observation that a couple of propositions still hold if “convex” is replaced by “quasi-convex”.

In this way we get a generalization of a theorem, where a *stronger* condition is replaced by a *weaker* one.

# Quasi-Convex and Quasi-Concave II

- ▶ Function  $f$  is *quasi-convex* if and only if


$$f((1 - h)\mathbf{x}_1 + h\mathbf{x}_2) \leq \max\{f(\mathbf{x}_1), f(\mathbf{x}_2)\}$$

for all  $\mathbf{x}_1, \mathbf{x}_2$  and  $h \in [0, 1]$ .


- ▶ Function  $f$  is *quasi-concave* if and only if

$$f((1 - h)\mathbf{x}_1 + h\mathbf{x}_2) \geq \min\{f(\mathbf{x}_1), f(\mathbf{x}_2)\}$$

for all  $\mathbf{x}_1, \mathbf{x}_2$  and  $h \in [0, 1]$ .



quasi-convex



quasi-concave

# Strictly Quasi-Convex and Quasi-Concave

- ▶ Function  $f$  is called **strictly quasi-convex** if

$$f((1 - h)\mathbf{x}_1 + h\mathbf{x}_2) < \max\{f(\mathbf{x}_1), f(\mathbf{x}_2)\}$$

for all  $\mathbf{x}_1, \mathbf{x}_2$ , with  $\mathbf{x}_1 \neq \mathbf{x}_2$ , and  $h \in (0, 1)$ .

- ▶ Function  $f$  is called **strictly quasi-concave** if

$$f((1 - h)\mathbf{x}_1 + h\mathbf{x}_2) > \min\{f(\mathbf{x}_1), f(\mathbf{x}_2)\}$$

for all  $\mathbf{x}_1, \mathbf{x}_2$ , with  $\mathbf{x}_1 \neq \mathbf{x}_2$ , and  $h \in (0, 1)$ .

# Quasi-convex and Quasi-Concave III

For a differentiable function  $f$  we find:

- ▶ Function  $f$  is *quasi-convex* if and only if

$$f(\mathbf{x}) \leq f(\mathbf{x}_0) \quad \Rightarrow \quad \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) \leq 0$$

- ▶ Function  $f$  is *quasi-concave* if and only if

$$f(\mathbf{x}) \geq f(\mathbf{x}_0) \quad \Rightarrow \quad \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) \geq 0$$

# Summary

- ▶ monotone function
- ▶ convex set
- ▶ convex and concave function
- ▶ convexity and definiteness of quadratic form
- ▶ minors of Hessian matrix
- ▶ quasi-convex and quasi-concave function