

## Provided functionality

- Bayesian estimation of finite mixtures of univariate Gaussian distributions with MCMC methods
- accompanying material to the book Frühwirth-Schnatter (2005)
- special purpose interface to JAGS (Just Another Gibbs Sampler; Plummer, 2004)

## Model class

Finite Gaussian mixture distributions are defined by

$$\mathbb{P}(y_i) = \sum_{k=1}^K \eta_k \text{Normal}(y_i | \mu_k, \sigma_k^2)$$

where  $\eta_k \geq 0$ ,  $\sum_{k=1}^K \eta_k = 1$  and  $\text{Normal}(\cdot | \mu, \sigma^2)$  is a Gaussian distribution with mean  $\mu$  and variance  $\sigma^2$ .

For the prior of  $\eta_k$  it is assumed that

$$\eta_1, \dots, \eta_K \sim \text{Dirichlet}(e_{0,1}, \dots, e_{0,K})$$

For conditionally conjugate priors we have

$$\sigma_k^2 \sim \text{InvGamma}\left(\frac{\nu_{0,k}}{2}, \frac{\nu_{0,k} S_{0,k}}{2}\right) \quad \mu_k | \sigma_k^2 \sim \text{Normal}(b_{0,k}, B_{0,k} \sigma_k^2)$$

For independence priors

$$\sigma_k^2 \sim \text{InvGamma}\left(\frac{\nu_{0,k}}{2}, \frac{\nu_{0,k} S_{0,k}}{2}\right) \quad \mu_k \sim \text{Normal}(b_{0,k}, B_{0,k})$$

For a hierarchical prior on the variances we add

$$S_{0,k} \sim \text{Gamma}\left(\frac{g_{0,k}}{2}, \frac{g_{0,k} G_{0,k}}{2}\right)$$

## Implementation

*Model specification:*

- number of segments
- specific prior distribution and parameter values
- initial values

*MCMC sampling:*

- number of burn-in and monitored draws
- random seed




Fig. 1: Flowchart between R and JAGS: R code necessary for creating input for JAGS in BUGS syntax and visualization of results in R

*Plot functions* for output analysis

- diagnostic plots: for determining a suitable ordering constraint and the number of segments (see Figure 2)
- output visualization: trace and density plots (see Figure 3)
- a-posteriori probabilities (see Figure 4)

## Application

"fish" dataset: (Titterton et al., 1985)

- 256 observations of fish length
- *unobserved heterogeneity* because of different age groups present in the population




Fig. 2: Diagnostic plots:  $\mu$  versus  $\sigma$  on the left and  $\mu_k$  versus  $\mu_l$  for  $k \neq l$  on the right




Fig. 3: Traces and density of the MCMC draws for  $\mu$  under the constraint  $\mu_1 < \dots < \mu_4$ .




Fig. 4: Posterior classification of the observations where the MCMC draws are sorted with respect to the constraint  $\mu_1 < \dots < \mu_4$ .

## References

- Frühwirth-Schnatter, S. 2005. *Bayesian Mixture Modelling*. Springer. Forthcoming.
- Plummer, M. 2004. *JAGS Version 0.50 manual*. International Agency for Research on Cancer.
- Titterton, D., Smith, A., & Makov, U. 1985. *Statistical Analysis of Finite Mixture Distributions*. Chichester: Wiley.